



Joint Pain Flares Among People With HIV, Hepatitis C or Both

May 5, 2015

People with HIV or hepatitis C virus (HCV), or who are coinfecting with both viruses, report high levels of joint pain, *aidsmap* reports. Publishing their findings in *BMC Musculoskeletal Disorders*, researchers conducted a cross-sectional study in which they interviewed 202 participants about joint pain, including 93 people with hep C, 30 people with HIV, and 79 HCV/HIV coinfecting people.

The groups had similar age and gender makeups. A total of 173 of the participants were male.

The researchers used the Multi-Dimensional Health Assessment Questionnaire to evaluate joint pain and associated symptoms.

Seventy-one percent of the HCV-monoinfecting people reported joint pain, compared with 56 percent of the HIV/HCV coinfecting people and 50 percent of the HIV-monoinfecting participants.

Across the study, the most commonly cited areas of joint pain were the fingers, knees and back.

Among the HCV-positive participants, having arthritis raised the risk of joint pain 4.25-fold, while smoking raised the risk 5.02-fold. Among the HIV-positive participants, arthritis raised the risk of joint pain 5.36-fold, and smoking raised the risk 6.07-fold.

To read the MedPage Today article, [click here](#).

To read the study, [click here](#).

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.hepmag.com/article/joint-pain-27167>