



Living With Liver Disease

Over-the-Counter Pain Remedies

Everything that starts in the mouth passes through the liver, including prescription, over-the-counter (OTC) and recreational drugs. Because of this, experts recommend that people with hepatitis and other liver diseases use caution when taking medications. A liver that doesn't have cirrhosis can usually metabolize medications that are taken as directed, assuming alcohol is not consumed around the same time. A cirrhotic liver may have difficulty metabolizing certain medications, thus your medical provider may reduce your medication dose.

Acetaminophen (Tylenol) is one of the most commonly used over-the-counter medications. In the U.S., approximately 50 million people take acetaminophen every week, and more than 25 billion doses are sold yearly. Known as paracetamol in Europe, acetaminophen is effective for headaches, fever and mild pain. It is harmless at low doses and one of the safest drugs there is, even for those with liver disease. However, acetaminophen can cause liver injury and death from acute liver failure at amounts just twice the maximum recommended dose of acetaminophen. Slightly more than 300 people die annually from overdose of the drug.

Roughly half of acetaminophen deaths are the result of suicide. The rest are unintentional overdoses. The big problem with acetaminophen is that it is added to many medications, including remedies for colds, headaches, pain, sleep, sinus problems, cough and menstrual discomfort. People may not know that they are taking acetaminophen, let alone too much of it.

The abbreviation "APAP" is often used on labels of drugs that have added acetaminophen. Be sure that your total APAP dose includes all sources of this drug. APAP is added to over 600 other medications, including pain meds, sleep meds, cold meds, cough meds, sinus meds, etc. You can check to see if a drug contains APAP here: www.getreliefresponsibly.com.

Excessive acetaminophen (Tylenol) is the leading cause of acute liver failure in the U.S. Despite research showing that low doses of acetaminophen is safe for most patients with liver disease, many people avoid taking it. If you are concerned about taking acetaminophen, talk to your doctor. All patients, especially those who are pre- and post- liver transplant, should ask their medical providers for a recommended dose. Here are some tips for taking acetaminophen

- Follow dosing instructions precisely. Never take more than 1,000 mg in a single dose.
- Start small. A single dose ranges from 325 mg, 500 mg, 650 mg or 1,000 mg; some people have good results at the lower doses.
- Do not take acetaminophen if you drink alcohol, especially if you have hepatitis or another liver disease.

Acetaminophen's reputation may lead you to switch to non-steroidal anti-inflammatory drugs (NSAIDs), such as aspirin, ibuprofen (Advil, Motrin, etc), and naproxen (Aleve). But regular use of NSAIDs at normal doses can cause stomach bleeding, kidney problems, and increase the chance of heart attack or stroke. Talk with your doctor about what type of pain relief will work best for you.

Last Reviewed: June 29, 2023

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.hepmag.com/basics/liver-health/over-the-counter-remedies>